

Low-lying quasibound states of H₂¹⁶O

T. Szidarovszky^a, A. G. Császár^b

^a Laboratory of Complex Chemical Systems, Tel.: (36 1) 372 2500 / 6505 , E-mail: tamas821@gmail.com

^b Laboratory of Molecular Structure and Dynamics, Eötvös Loránd University, H-1117, Budapest, Pázmány Péter sétány 1/A. Tel.: (36 1) 372-2929 , Fax: (36 1) 372-2592 E-mail:

Quasibound (also called resonance) states of a system are metastable states which have sufficient energy to brake up the system into its subsystems. They can be associated with outgoing eigenfunctions of the Hamiltonian, diverging exponentially at infinity. Due to the outgoing boundary condition, the eigenvalues corresponding to resonance states are complex.

Techniques of non-Hermitian quantum chemistry merged with efficient nuclear motion computational algorithms allow the determination of quasibound rovibrational states of strongly-bound triatomic molecules.

The molecule chosen to present the methods is H₂¹⁶O, for which an accurate global potential energy surface [1], previous computation of resonance states [2], and some Feshbach ($J = 0$, where J is the quantum number characterizing overall rotations of the molecule) and shape ($J \neq 0$) resonances measured via a state-selective, triple-resonance technique [3] are available.

Characterization of the computed resonance states is performed via probability density plots (see Fig. 1.) based on rovibrational wavefunctions obtained as the eigenfunctions of non-Hermitian Hamiltonians. Just like for bound states, such plots provide useful details about the nature of the resonance states, such as dissociative behaviour, effect of rotational excitation on resonance state lifetimes, etc..

Interested readers can find more in depth information in a recently published paper [4], which covers most of the presented results.




Figure 1. Two-dimensional sections of probability density plots obtained from eigenfunctions of non-Hermitian rovibrational Hamiltonians, where R_1 and R_2 are the two OH bond lengths and θ is the HOH bond angle. The area near the equilibrium structure is also shown enlarged in the leftmost picture.

- [1] A. G. Császár, E. Mátyus, T. Szidarovszky, L. Lodi, N. F. Zobov, S. V. Shirin, O. L. Polyansky and J. Tennyson, *J. Quant. Spectr. Rad. Transfer*, (**2010**), 111, 1043
- [2] N. F. Zobov, S. V. Shirin, L. Lodi, B. C. Silva, J. Tennyson, A. G. Császár and O. L. Polyansky, *Chem. Phys. Lett.*, (**2011**), 507, 48
- [3] M. Grechko, P. Maksyutenko, T. R. Rizzo and O. V. Boyarkin, *J. Chem. Phys.*, (**2010**), 133, 081103
- [4] T. Szidarovszky, A. G. Császár, *Mol. Phys. (Martin Quack Special Issue)*, (**2013**), 111, 2113